

## Chapter 5


# Simple applications of macroscopic thermodynamics

Speaker: Yufeng Lu  
Supervisor: Dr. Jun Fan

City University of Hong Kong

Oct. 17, 2014

# Line



# Line

Macro parameters ( $E, T...$ )



Thermodynamic laws

Basic relations ( $TdS = dE + pdV...$ )



Thermodynamic functions

Maxwell relations



General relations ( $C_p - C_V$  from  $V, T, p...$ )

Why the last step?

experimental measurements  $\rightarrow$  theoretical properties.

# Outline

- 1 **An example:**  $TdS = dE + pdV$ .
- 2 **A special case: for ideal gases**
  - Energy:  $E = E(T)$
  - Heat capacities:  $c_p = c_V + R$
  - Quasi-static adiabatic process:  $pV^\gamma = \text{constant}$
- 3 **Maxwell relations and thermodynamic functions**
- 4 **General cases**
  - Heat capacities, after Maxwell relations
  - Energy, after Maxwell relations
- 5 **Applications**
  - Heat engines
  - Refrigerators
  - Equivalence of Kelvin and Clausius statements

# Outline

- 1 **An example:**  $TdS = dE + pdV$ .
- 2 **A special case: for ideal gases**
  - Energy:  $E = E(T)$
  - Heat capacities:  $c_p = c_V + R$
  - Quasi-static adiabatic process:  $pV^\gamma = \text{constant}$
- 3 **Maxwell relations and thermodynamic functions**
- 4 **General cases**
  - Heat capacities, after Maxwell relations
  - Energy, after Maxwell relations
- 5 **Applications**
  - Heat engines
  - Refrigerators
  - Equivalence of Kelvin and Clausius statements

# An example

An example..

# An example

An example..

Examples

## An example

An example..

Examples are always **useful**.



## An example

To an infinitesimal process,

## An example

To an **infinitesimal process**, the first law yields

## An example

To an **infinitesimal process**, the first law yields

$$dQ = dE + dW. \quad (1.1)$$

## An example

To an **infinitesimal process**, the first law yields

$$dQ = dE + dW. \quad (1.1)$$

If the process is also **quasi-static**,

## An example

To an **infinitesimal process**, the first law yields

$$dQ = dE + dW. \quad (1.1)$$

If the process is also **quasi-static**, the second law says

$$dQ = TdS. \quad (1.2)$$

## An example

To an **infinitesimal process**, the first law yields

$$dQ = dE + dW. \quad (1.1)$$

If the process is also **quasi-static**, the second law says

$$dQ = TdS. \quad (1.2)$$

As well known,

## An example

To an **infinitesimal process**, the first law yields

$$dQ = dE + dW. \quad (1.1)$$

If the process is also **quasi-static**, the second law says

$$dQ = TdS. \quad (1.2)$$

As well known, the work done **by** the system is given by

$$dW = pdV. \quad (1.3)$$

## An example

To an **infinitesimal process**, the first law yields

$$dQ = dE + dW. \quad (1.1)$$

If the process is also **quasi-static**, the second law says

$$dQ = TdS. \quad (1.2)$$

As well known, the work done **by** the system is given by

$$dW = pdV. \quad (1.3)$$

Combining (1.1) to (1.3), one may get that

$$TdS = dE + pdV. \quad (5.1.5)$$



## The aim of this chapter

The whole chapter will be based on

## The aim of this chapter

The whole chapter will be based on **only** the thermodynamic laws

## The aim of this chapter

The whole chapter will be based on **only** the thermodynamic laws to derive important **relationships** between **macroscopic** quantities.

## The aim of this chapter

The whole chapter will be based on **only** the thermodynamic laws to derive important **relationships** between **macroscopic** quantities. Those conclusions are **completely independent** of

## The aim of this chapter

The whole chapter will be based on **only** the thermodynamic laws to derive important **relationships** between **macroscopic** quantities. Those conclusions are **completely independent** of any specific models assumed to describe

## The aim of this chapter

The whole chapter will be based on **only** the thermodynamic laws to derive important **relationships** between **macroscopic** quantities. Those conclusions are **completely independent** of any specific models assumed to describe the **microscopic** constituents of a system.

# Outline

- 1 An example:  $TdS = dE + pdV$ .
- 2 **A special case: for ideal gases**
  - Energy:  $E = E(T)$
  - Heat capacities:  $c_p = c_V + R$
  - Quasi-static adiabatic process:  $pV^\gamma = \text{constant}$
- 3 Maxwell relations and thermodynamic functions
- 4 General cases
  - Heat capacities, after Maxwell relations
  - Energy, after Maxwell relations
- 5 Applications
  - Heat engines
  - Refrigerators
  - Equivalence of Kelvin and Clausius statements

# Energy



# Energy

## Theorem

*For an ideal gas, its internal energy does not depend on its volume, but only on its temperature, which is*

$$E = E(T).$$

# Energy

## Theorem

*For an ideal gas, its internal energy does not depend on its volume, but only on its temperature, which is*

$$E = E(T).$$

*proof.* Suppose that there are  $\nu$  mole ideal gas.

# Energy

## Theorem

*For an ideal gas, its internal energy does not depend on its volume, but only on its temperature, which is*

$$E = E(T).$$

*proof.* Suppose that there are  $\nu$  mole ideal gas. The [equation of states](#) and (5.1.5) tell us,

$$\left. \begin{aligned} T dS &= dE + p dV; \\ \frac{p}{T} &= \frac{\nu R}{V}; \end{aligned} \right\} \Rightarrow dS = \frac{1}{T} dE + \frac{\nu R}{V} dV. \quad (2.1)$$

# Energy

## Theorem

*For an ideal gas, its internal energy does not depend on its volume, but only on its temperature, which is*

$$E = E(T).$$

*proof.* Suppose that there are  $\nu$  mole ideal gas. The [equation of states](#) and (5.1.5) tell us,

$$\left. \begin{aligned} T dS &= dE + p dV; \\ \frac{p}{T} &= \frac{\nu R}{V}; \end{aligned} \right\} \Rightarrow dS = \frac{1}{T} dE + \frac{\nu R}{V} dV. \quad (2.1)$$

From [the exact differential of  \$E\$](#) , one may obtain that

$$dE = \left( \frac{\partial E}{\partial T} \right)_V dT + \left( \frac{\partial E}{\partial V} \right)_T dV. \quad (2.2)$$

# Energy

Combining (2.1) and (2.2), one gets that

# Energy

Combining (2.1) and (2.2), one gets that

$$dS = \frac{1}{T} \left( \frac{\partial E}{\partial T} \right)_V dT + \left[ \frac{\nu R}{V} + \left( \frac{\partial E}{\partial V} \right)_T \right] dV.$$

# Energy

Combining (2.1) and (2.2), one gets that

$$dS = \frac{1}{T} \left( \frac{\partial E}{\partial T} \right)_V dT + \left[ \frac{\nu R}{V} + \left( \frac{\partial E}{\partial V} \right)_T \right] dV.$$

Compared with

$$dS = \left( \frac{\partial S}{\partial T} \right)_V dT + \left( \frac{\partial S}{\partial V} \right)_T dV,$$

# Energy

Combining (2.1) and (2.2), one gets that

$$dS = \frac{1}{T} \left( \frac{\partial E}{\partial T} \right)_V dT + \left[ \frac{\nu R}{V} + \left( \frac{\partial E}{\partial V} \right)_T \right] dV.$$

Compared with

$$dS = \left( \frac{\partial S}{\partial T} \right)_V dT + \left( \frac{\partial S}{\partial V} \right)_T dV,$$

one has

$$\begin{aligned} \left( \frac{\partial S}{\partial T} \right)_V &= \frac{1}{T} \left( \frac{\partial E}{\partial T} \right)_V; \\ \left( \frac{\partial S}{\partial V} \right)_T &= \frac{1}{T} \left( \frac{\partial E}{\partial V} \right)_T + \frac{\nu R}{V}. \end{aligned}$$



# Energy

Combining (2.1) and (2.2), one gets that

$$dS = \frac{1}{T} \left( \frac{\partial E}{\partial T} \right)_V dT + \left[ \frac{\nu R}{V} + \left( \frac{\partial E}{\partial V} \right)_T \right] dV.$$

Compared with

$$dS = \left( \frac{\partial S}{\partial T} \right)_V dT + \left( \frac{\partial S}{\partial V} \right)_T dV,$$

one has

$$\begin{aligned} \left( \frac{\partial}{\partial V} \right)_T \left( \frac{\partial S}{\partial T} \right)_V &= \left( \frac{\partial}{\partial V} \right)_T \left[ \frac{1}{T} \left( \frac{\partial E}{\partial T} \right)_V \right]; \\ \left( \frac{\partial}{\partial T} \right)_V \left( \frac{\partial S}{\partial V} \right)_T &= \left( \frac{\partial}{\partial T} \right)_V \left[ \frac{1}{T} \left( \frac{\partial E}{\partial V} \right)_T + \frac{\nu R}{V} \right]. \end{aligned}$$

# Energy

By the **equality of the second derivatives** for  $S(T, V)$ ,

# Energy

By the **equality of the second derivatives** for  $S(T, V)$ ,

$$\left(\frac{\partial}{\partial V}\right)_T \left[ \frac{1}{T} \left(\frac{\partial E}{\partial T}\right)_V \right] = \left(\frac{\partial}{\partial T}\right)_V \left[ \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T + \frac{\nu R}{V} \right],$$

# Energy

By the **equality of the second derivatives** for  $S(T, V)$ ,

$$\left(\frac{\partial}{\partial V}\right)_T \left[ \frac{1}{T} \left(\frac{\partial E}{\partial T}\right)_V \right] = \left(\frac{\partial}{\partial T}\right)_V \left[ \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T + \frac{\nu R}{V} \right],$$

which further implies

$$\frac{1}{T} \cancel{\frac{\partial^2 E}{\partial V \partial T}} = \left(\frac{\partial}{\partial T}\right)_V \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T + \frac{1}{T} \cancel{\frac{\partial^2 E}{\partial T \partial V}} + 0.$$

# Energy

By the **equality of the second derivatives** for  $S(T, V)$ ,

$$\left(\frac{\partial}{\partial V}\right)_T \left[ \frac{1}{T} \left(\frac{\partial E}{\partial T}\right)_V \right] = \left(\frac{\partial}{\partial T}\right)_V \left[ \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T + \frac{\nu R}{V} \right],$$

which further implies

$$= \left(\frac{\partial}{\partial T}\right)_V \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T .$$

# Energy

By the **equality of the second derivatives** for  $S(T, V)$ ,

$$\left(\frac{\partial}{\partial V}\right)_T \left[ \frac{1}{T} \left(\frac{\partial E}{\partial T}\right)_V \right] = \left(\frac{\partial}{\partial T}\right)_V \left[ \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T + \frac{\nu R}{V} \right],$$

which further implies

$$= \left(\frac{\partial}{\partial T}\right)_V \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T .$$

Therefore,

$$\left(\frac{\partial E}{\partial V}\right)_T = 0,$$

# Energy

By the **equality of the second derivatives** for  $S(T, V)$ ,

$$\left(\frac{\partial}{\partial V}\right)_T \left[ \frac{1}{T} \left(\frac{\partial E}{\partial T}\right)_V \right] = \left(\frac{\partial}{\partial T}\right)_V \left[ \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T + \frac{\nu R}{V} \right],$$

which further implies

$$= \left(\frac{\partial}{\partial T}\right)_V \frac{1}{T} \left(\frac{\partial E}{\partial V}\right)_T .$$

Therefore,

$$\left(\frac{\partial E}{\partial V}\right)_T = 0,$$

which means the internal energy of an ideal gas is not dependent on its volume.

# Heat capacities



# Heat capacities

Consider a system  $A = A(T, y)$ .

# Heat capacities

Consider a system  $A = A(T, y)$ .  $y$  probably be  $V$  or  $p$ .

# Heat capacities

Consider a system  $A = A(T, y)$ .  $y$  probably be  $V$  or  $p$ . Recall that,

$$C_y := \left( \frac{dQ}{dT} \right)_y, \quad (2.3)$$

and

# Heat capacities

Consider a system  $A = A(T, y)$ .  $y$  probably be  $V$  or  $p$ . Recall that,

$$C_y := \left( \frac{dQ}{dT} \right)_y, \quad (2.3)$$

and

$$c_y := \frac{1}{\nu} C_y = \frac{1}{\nu} \left( \frac{dQ}{dT} \right)_y. \quad (2.4)$$

# Heat capacities

In an infinitesimal process,

# Heat capacities

In an **infinitesimal process**, the first law tells us

$$dQ = dE + pdV.$$

# Heat capacities

In an **infinitesimal process**, the first law tells us

$$dQ = dE + pdV.$$

In **any case**, for  $\nu$  mole an **ideal gas**, (should be in an **infinitesimal process**) by the definition of *heat capacity with constant volume*,

$$\nu c_V = C_V := \left( \frac{dQ}{dT} \right)_V = \frac{dE}{dT}.$$

# Heat capacities

In an **infinitesimal process**, the first law tells us

$$dQ = dE + pdV.$$

In **any case**, for  $\nu$  mole an **ideal gas**, (should be in an **infinitesimal process**) by the definition of *heat capacity with constant volume*,

$$\nu c_V = C_V := \left( \frac{dQ}{dT} \right)_V = \frac{dE}{dT}.$$

Thus,

$$dE = \nu c_V dT.$$



## Heat capacities

Similar, in **any case**, for an ideal gas, by the definition of *heat capacity with constant pressure*,

$$\begin{aligned}\nu c_p &:= \left( \frac{dQ}{dT} \right)_p = \left( \frac{dE + pdV}{dT} \right)_p \\ &= \left( \frac{dE}{dT} \right)_p + \left( \frac{pdV}{dT} \right)_p\end{aligned}$$

## Heat capacities

Similar, in **any case**, for an ideal gas, by the definition of *heat capacity with constant pressure*,

$$\begin{aligned}\nu c_p &:= \left( \frac{dQ}{dT} \right)_p = \left( \frac{dE + pdV}{dT} \right)_p \\ &= \left( \frac{dE}{dT} \right)_p + \left( \frac{pdV}{dT} \right)_p = \nu c_V + \nu R.\end{aligned}$$

## Heat capacities

Similar, in **any case**, for an ideal gas, by the definition of *heat capacity with constant pressure*,

$$\begin{aligned}\nu c_p &:= \left( \frac{dQ}{dT} \right)_p = \left( \frac{dE + pdV}{dT} \right)_p \\ &= \left( \frac{dE}{dT} \right)_p + \left( \frac{pdV}{dT} \right)_p = \nu c_V + \nu R.\end{aligned}$$

Thus,

$$c_p = c_V + R.$$

# Heat capacities

Similar, in **any case**, for an ideal gas, by the definition of *heat capacity with constant pressure*,

$$\begin{aligned}\nu c_p &:= \left( \frac{dQ}{dT} \right)_p = \left( \frac{dE + pdV}{dT} \right)_p \\ &= \left( \frac{dE}{dT} \right)_p + \left( \frac{pdV}{dT} \right)_p = \nu c_V + \nu R.\end{aligned}$$

Thus,

$$c_p = c_V + R.$$

$c_p \rightarrow$  more easily to get by **experiment**.

## Heat capacities

Similar, in **any case**, for an ideal gas, by the definition of *heat capacity with constant pressure*,

$$\begin{aligned}\nu c_p &:= \left( \frac{dQ}{dT} \right)_p = \left( \frac{dE + pdV}{dT} \right)_p \\ &= \left( \frac{dE}{dT} \right)_p + \left( \frac{pdV}{dT} \right)_p = \nu c_V + \nu R.\end{aligned}$$

Thus,

$$c_p = c_V + R.$$

$c_p \rightarrow$  more easily to get by **experiment**.

$c_V \rightarrow$  more easily to get by **theoretic calculation**.

# Quasi-static adiabatic process

## Quasi-static adiabatic process

- For an ideal gas,  $T$  is keeping constant  $\Rightarrow$

$$pV = \text{const.} \Rightarrow p \downarrow = \frac{\text{const.}}{V \uparrow}$$

decreases when the gas expands.

## Quasi-static adiabatic process

- For an ideal gas,  $T$  is keeping constant  $\Rightarrow$

$$pV = \text{const.} \Rightarrow p \downarrow = \frac{\text{const.}}{V \uparrow}$$

decreases when the gas expands.

- For an ideal gas, when it expands under an adiabatic ( $dQ = 0$ ) condition,

$$0 = dQ = dE \downarrow + dW \uparrow \Rightarrow T \downarrow = 2E \downarrow / 3Nk.$$



## Quasi-static adiabatic process

- For an ideal gas,  $T$  is keeping constant  $\Rightarrow$

$$pV = \text{const.} \Rightarrow p \downarrow = \frac{\text{const.}}{V \uparrow}$$

decreases when the gas expands.

- For an ideal gas, when it expands under an adiabatic ( $dQ = 0$ ) condition,

$$0 = dQ = dE \downarrow + dW \uparrow \Rightarrow T \downarrow = 2E \downarrow / 3Nk.$$

- Question: How is the pressure  $p$  related to  $V$  in a kind of the second process?

## Quasi-static adiabatic process


### Theorem

*In the second process,*

$$pV^\gamma = \text{const.} \quad (2.5)$$

where  $\gamma := \frac{c_p}{c_v} > 1$ .

# Quasi-static adiabatic process



# Outline

- 1 An example:  $TdS = dE + pdV$ .
- 2 A special case: for ideal gases
  - Energy:  $E = E(T)$
  - Heat capacities:  $c_p = c_V + R$
  - Quasi-static adiabatic process:  $pV^\gamma = \text{constant}$
- 3 **Maxwell relations and thermodynamic functions**
- 4 **General cases**
  - Heat capacities, after Maxwell relations
  - Energy, after Maxwell relations
- 5 **Applications**
  - Heat engines
  - Refrigerators
  - Equivalence of Kelvin and Clausius statements

# Maxwell relations

Consider a quasi-static infinitesimal process with a system

## Maxwell relations

Consider a quasi-static infinitesimal process with a system having only  $V$  as an external parameter.

# Maxwell relations

Consider a quasi-static infinitesimal process with a system having only  $V$  as an external parameter. one should have

$$dQ = TdS = dE + pdV. \quad (3.1)$$

$$E = E(S, V)$$

From (3.1), one can find  $dE = TdS - pdV$ .



$$E = E(S, V)$$

From (3.1), one can find  $dE = TdS - pdV$ . Further,

$$E = E(S, V)$$

From (3.1), one can find  $dE = TdS - pdV$ . Further,

$$\left. \begin{aligned} \left( \frac{\partial E}{\partial S} \right)_V &= T(S, V); \\ \left( \frac{\partial E}{\partial V} \right)_S &= -p(S, V); \\ \frac{\partial^2 E}{\partial V \partial S} &= \frac{\partial^2 E}{\partial S \partial V} \end{aligned} \right\} \quad (3.2)$$

$$E = E(S, V)$$

From (3.1), one can find  $dE = TdS - pdV$ . Further,

$$\left. \begin{aligned} \left( \frac{\partial E}{\partial S} \right)_V &= T(S, V); \\ \left( \frac{\partial E}{\partial V} \right)_S &= -p(S, V); \\ \frac{\partial^2 E}{\partial V \partial S} &= \frac{\partial^2 E}{\partial S \partial V} \end{aligned} \right\} \Rightarrow \left( \frac{\partial T}{\partial V} \right)_S = - \left( \frac{\partial p}{\partial S} \right)_V. \quad (3.2)$$

$$E = E(S, V)$$

From (3.1), one can find  $dE = TdS - pdV$ . Further,

$$\left. \begin{aligned} \left( \frac{\partial E}{\partial S} \right)_V &= T(S, V); \\ \left( \frac{\partial E}{\partial V} \right)_S &= -p(S, V); \\ \frac{\partial^2 E}{\partial V \partial S} &= \frac{\partial^2 E}{\partial S \partial V} \end{aligned} \right\} \Rightarrow \left( \frac{\partial T}{\partial V} \right)_S = - \left( \frac{\partial p}{\partial S} \right)_V. \quad (3.2)$$

**Note:** here,  $dE$  is used as an exact differential.

$$H = H(S, p)$$

Again, consider the same process and the same system as previous.

$$H = H(S, p)$$

Again, consider the same process and the same system as previous. Let  $V = V(p)$ . Then

$$\frac{d(pV(p))}{dp} = V(p) + p \frac{dV(p)}{dp}.$$

Equivalently,

$$pdV = d(pV) - Vdp.$$

$$H = H(S, p)$$

Again, consider the same process and the same system as previous. Let  $V = V(p)$ . Then

$$\frac{d(pV(p))}{dp} = V(p) + p \frac{dV(p)}{dp}.$$

Equivalently,

$$pdV = d(pV) - Vdp.$$

From again (3.1),

$$H = H(S, p)$$

Again, consider the same process and the same system as previous. Let  $V = V(p)$ . Then

$$\frac{d(pV(p))}{dp} = V(p) + p \frac{dV(p)}{dp}.$$

Equivalently,

$$pdV = d(pV) - Vdp.$$

From again (3.1), i.e.  $TdS = dE + pdV$ ,



$$H = H(S, p)$$

Again, consider the same process and the same system as previous. Let  $V = V(p)$ . Then

$$\frac{d(pV(p))}{dp} = V(p) + p \frac{V(p)}{dp}.$$

Equivalently,

$$pdV = d(pV) - Vdp.$$

From again (3.1), i.e.  $TdS = dE + pdV$ , one has

$$\begin{aligned} TdS &= dE + d(pV) - Vdp \\ \Rightarrow d(E + pV) &= TdS + Vdp. \end{aligned} \tag{3.3}$$

$$H = H(S, p)$$

Therefore, one can find  $d(E + pV)$  is another exact differential.

$$H = H(S, p)$$

Therefore, one can find  $d(E + pV)$  is another exact differential.  
Define

$$H := E + pV$$

and call it

$$H = H(S, p)$$

Therefore, one can find  $d(E + pV)$  is another exact differential.  
Define

$$H := E + pV$$

and call it “enthalpy”.

$$H = H(S, p)$$

Therefore, one can find  $d(E + pV)$  is another exact differential.  
Define

$$H := E + pV$$

and call it “**enthalpy**”. As you can imagine,

$$H = H(S, p)$$

Therefore, one can find  $d(E + pV)$  is another exact differential.  
Define

$$H := E + pV$$

and call it “**enthalpy**”. As you can imagine, **something should be done to  $H$**  as treated to  $E$  before.

$$H = H(S, p)$$

By **definition** of  $H$ , one has  $dH = TdS + Vdp$  and further,

$$H = H(S, p)$$

By **definition** of  $H$ , one has  $dH = TdS + Vdp$  and further,

$$\left. \begin{aligned} \left( \frac{\partial H}{\partial S} \right)_p &= T(S, p); \\ \left( \frac{\partial H}{\partial p} \right)_S &= V(S, p); \\ \frac{\partial^2 H}{\partial p \partial S} &= \frac{\partial^2 H}{\partial S \partial p} \end{aligned} \right\} \quad \left( \frac{\partial T}{\partial p} \right)_S = - \left( \frac{\partial V}{\partial S} \right)_p. \quad (3.4)$$



$$F = F(T, V)$$

Again and again, consider the same system and process as before.

$$F = F(T, V)$$

Again and again, consider the same system and process as before. Very similarly, one can find  $d(E - TS)$  should be an exact differential. So define a quantity

$$F := E - TS,$$

where  $E$  and  $S$  are treated as functions of  $T$  and  $V$ .

$$F = F(T, V)$$

Again and again, consider the same system and process as before. Very similarly, one can find  $d(E - TS)$  should be an exact differential. So define a quantity

$$F := E - TS,$$

where  $E$  and  $S$  are treated as functions of  $T$  and  $V$ .  $F$  then is usually called the “Helmholtz free energy”.

$$F = F(T, V)$$

Again and again, consider the **same** system and process as before. Very similarly, one can find  $d(E - TS)$  should be an exact differential. So define a quantity

$$F := E - TS,$$

where  $E$  and  $S$  are treated as functions of  $T$  and  $V$ .

$F$  then is usually called the “**Helmholtz free energy**”.

And, by using the equality of the cross derivatives, one may obtain that

$$\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial p}{\partial T}\right)_V. \quad (3.5)$$

$$G = G(T, p)$$

Again and again and again, the same system and process are under considered.

$$G = G(T, p)$$

Again and again and again, the same system and process are under considered. One can also find  $d(E - TS + pV)$  is another exact differential. Define

$$G := E - TS + pV$$

and call it “Gibbs free energy”.

$$G = G(T, p)$$

Again and again and again, the same system and process are under considered. One can also find  $d(E - TS + pV)$  is another exact differential. Define

$$G := E - TS + pV$$

and call it “Gibbs free energy”. Again, the equality of the cross derivatives tells us

$$-\left(\frac{\partial S}{\partial p}\right)_T = \left(\frac{\partial V}{\partial T}\right)_p. \quad (3.6)$$

# Summarize

## Maxwell relations

$$\left(\frac{\partial T}{\partial V}\right)_S = -\left(\frac{\partial p}{\partial S}\right)_V$$

$$\left(\frac{\partial T}{\partial p}\right)_S = -\left(\frac{\partial V}{\partial S}\right)_p$$

$$\left(\frac{\partial S}{\partial V}\right)_T = \left(\frac{\partial p}{\partial T}\right)_V$$

$$-\left(\frac{\partial S}{\partial p}\right)_T = \left(\frac{\partial V}{\partial T}\right)_p$$

## Thermodynamic functions

$$dE = TdS - pdV$$

$$dH = TdS + Vdp$$

$$dF = -SdT - pdV$$

$$dG = -SdT + Vdp$$



# Outline

- 1 An example:  $TdS = dE + pdV$ .
- 2 A special case: for ideal gases
  - Energy:  $E = E(T)$
  - Heat capacities:  $c_p = c_V + R$
  - Quasi-static adiabatic process:  $pV^\gamma = \text{constant}$
- 3 Maxwell relations and thermodynamic functions
- 4 General cases
  - Heat capacities, after Maxwell relations
  - Energy, after Maxwell relations
- 5 Applications
  - Heat engines
  - Refrigerators
  - Equivalence of Kelvin and Clausius statements

## Heat capacities

One can use the [Maxwell relations](#) for some substance to deduce the relation between the [heat capacity under constant volume](#) and the [heat capacity under constant pressure](#).

## Heat capacities

One can use the [Maxwell relations](#) for [some substance](#) to deduce the relation between the [heat capacity under constant volume](#) and the [heat capacity under constant pressure](#).

### Theorem

Let

$$\alpha := \frac{1}{V} \left( \frac{\partial V}{\partial T} \right)_p ;$$

$$\kappa := -\frac{1}{V} \left( \frac{\partial V}{\partial p} \right)_T .$$

Then,

$$C_p - C_V = VT \frac{\alpha^2}{\kappa} . \quad (4.1)$$

# Energy

One can also use the [Maxwell relations](#) to deduce the [internal energy](#) of [some substance](#) as a function of volume and temperature.

# Energy

One can also use the [Maxwell relations](#) to deduce the [internal energy](#) of [some substance](#) as a function of volume and temperature.

## Theorem

*Treat  $E$  as a function of  $V$  and  $T$ . Then,*


$$\left(\frac{\partial E}{\partial T}\right)_V = C_V;$$

$$\left(\frac{\partial E}{\partial V}\right)_T = T \left(\frac{\partial p}{\partial T}\right)_V - p.$$

# Outline


- 1 An example:  $TdS = dE + pdV$ .
- 2 A special case: for ideal gases
  - Energy:  $E = E(T)$
  - Heat capacities:  $c_p = c_V + R$
  - Quasi-static adiabatic process:  $pV^\gamma = \text{constant}$
- 3 Maxwell relations and thermodynamic functions
- 4 General cases
  - Heat capacities, after Maxwell relations
  - Energy, after Maxwell relations
- 5 Applications
  - Heat engines
  - Refrigerators
  - Equivalence of Kelvin and Clausius statements

# What is a heat engine?



$$T_1 > T_2$$

# What is a heat engine?




Energy conservation  $\Rightarrow$

$$Q_1 = W + Q_2.$$



# What is a heat engine?




Energy conservation  $\Rightarrow$

$$Q_1 = W + Q_2.$$

Define

$$\eta := \frac{W}{Q_1} \leq 1.$$

# What is a heat engine?



Energy conservation  $\Rightarrow$


$$Q_1 = W + Q_2.$$

Define

$$\eta := \frac{W}{Q_1} \leq 1.$$

$$\eta = 1 \Rightarrow W = Q_1$$

# What is a heat engine?



Energy conservation  $\Rightarrow$

$$Q_1 = W + Q_2.$$


Define

$$\eta := \frac{W}{Q_1} \leq 1.$$

$$\eta = 1 \Rightarrow W = Q_1$$


Perfect!

# Kelvin statement



Kelvin: Impossible to construct a device, operating in circle, that produce no effect other than the extraction of heat from a reservoir and do equivalent amount of work. <sup>1</sup>

# Kelvin statement



Kelvin: Impossible to construct a device, operating in circle, that produce no effect other than the extraction of heat from a reservoir and do equivalent amount of work. <sup>1</sup>

---

<sup>1</sup>P. Jacobs, *Thermodynamics*, Imperial College Press, 2013.

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $W > 0$ .

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $W > 0$ .
- Perfect requires  $Q_1 = W$ .

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $W > 0$ .
- Perfect requires  $Q_1 = W$ .
- Circle means **no entropy change to the device.**



## Why not?

- Remember  $Q_1 = W + Q_2$  and  $W > 0$ .
- Perfect requires  $Q_1 = W$ .
- Circle means no entropy change to the device.
- No effect means no entropy change to the outside.

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $W > 0$ .
- Perfect requires  $Q_1 = W$ .
- Circle means **no entropy change to the device**.
- No effect means **no entropy change to the outside**.
- So that

$$\Delta S = -\frac{\Delta Q_1}{T_1} = -\frac{W}{T_1} < 0,$$


## Why not?

- Remember  $Q_1 = W + Q_2$  and  $W > 0$ .
- Perfect requires  $Q_1 = W$ .
- Circle means **no entropy change to the device.**
- No effect means **no entropy change to the outside.**
- So that

$$\Delta S = -\frac{\Delta Q_1}{T_1} = -\frac{W}{T_1} < 0,$$


which is a contradiction with the second law.

# What is a refrigerator?



$$T_1 > T_2$$


# What is a refrigerator?



Energy conservation  $\Rightarrow$

$$Q_1 = W + Q_2.$$

# What is a refrigerator?




Energy conservation  $\Rightarrow$

$$Q_1 = W + Q_2.$$

Define

$$\eta' := \frac{Q_2}{Q_1} \leq 1.$$

# What is a refrigerator?



Energy conservation  $\Rightarrow$


$$Q_1 = W + Q_2.$$

Define

$$\eta' := \frac{Q_2}{Q_1} \leq 1.$$

$$\eta' = 1 \Rightarrow Q_2 = Q_1$$

# What is a refrigerator?



Energy conservation  $\Rightarrow$

$$Q_1 = W + Q_2.$$

Define

$$\eta' := \frac{Q_2}{Q_1} \leq 1.$$

$$\eta' = 1 \Rightarrow Q_2 = Q_1$$

Perfect!



# Clausius statement



Clausius: **Impossible** to construct a device, operating in circle, that produce **no effect** other than flowing **heat from a colder to a hotter body**.<sup>2</sup>

---

<sup>2</sup>P. Jacobs, *Thermodynamics*, Imperial College Press, 2013.

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $T_1 > T_2$ .

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $T_1 > T_2$ .
- Perfect requires  $Q_1 = Q_2$ .

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $T_1 > T_2$ .
- Perfect requires  $Q_1 = Q_2$ .
- Circle means **no entropy change to the device.**

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $T_1 > T_2$ .
- Perfect requires  $Q_1 = Q_2$ .
- Circle means **no entropy change to the device**.
- No effect means **no entropy change to the outside**.

## Why not?

- Remember  $Q_1 = W + Q_2$  and  $T_1 > T_2$ .
- Perfect requires  $Q_1 = Q_2$ .
- Circle means **no entropy change to the device.**
- No effect means **no entropy change to the outside.**
- So that

$$\Delta S = \frac{\Delta Q_1}{T_1} - \frac{\Delta Q_2}{T_2} = Q_1 \left( \frac{1}{T_1} - \frac{1}{T_2} \right) < 0,$$


## Why not?

- Remember  $Q_1 = W + Q_2$  and  $T_1 > T_2$ .
- Perfect requires  $Q_1 = Q_2$ .
- Circle means **no entropy change to the device**.
- No effect means **no entropy change to the outside**.
- So that

$$\Delta S = \frac{\Delta Q_1}{T_1} - \frac{\Delta Q_2}{T_2} = Q_1 \left( \frac{1}{T_1} - \frac{1}{T_2} \right) < 0,$$

which is a contradiction with the second law.


# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:




# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:

- Kelvin is wrong  $\Rightarrow$   
 $M_1$  possible ( $W = Q$ ).


# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:

- Kelvin is wrong  $\Rightarrow$   $M_1$  possible ( $W = Q$ ).
- For  $M_2$ ,  $\Rightarrow$   $Q_2 = W + Q_1 = Q + Q_1$ .


# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:

- Kelvin is wrong  $\Rightarrow$   $M_1$  possible ( $W = Q$ ).
- For  $M_2$ ,  $\Rightarrow$   $Q_2 = W + Q_1 = Q + Q_1$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1$  from  $T_2$ ;


# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:

- Kelvin is wrong  $\Rightarrow$   $M_1$  possible ( $W = Q$ ).
- For  $M_2$ ,  $\Rightarrow$   $Q_2 = W + Q_1 = Q + Q_1$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1$  from  $T_2$ ;
- while to eject  $Q_2 - Q = Q_1$  to  $T_1$ ;


# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:

- Kelvin is wrong  $\Rightarrow$   $M_1$  possible ( $W = Q$ ).
- For  $M_2$ ,  $\Rightarrow$   $Q_2 = W + Q_1 = Q + Q_1$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1$  from  $T_2$ ;
- while to eject  $Q_2 - Q = Q_1$  to  $T_1$ ;
- Remember  $T_1 > T_2$ .


# Equivalence



“Clausius  $\Rightarrow$  Kelvin”:


- Kelvin is wrong  $\Rightarrow$   $M_1$  possible ( $W = Q$ ).
- For  $M_2$ ,  $\Rightarrow$   $Q_2 = W + Q_1 = Q + Q_1$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1$  from  $T_2$ ;
- while to eject  $Q_2 - Q = Q_1$  to  $T_1$ ;
- Remember  $T_1 > T_2$ .
- This contradicts with Clausius statement.

# Equivalence



“Kelvin  $\Rightarrow$  Clausius”:

# Equivalence




“Kelvin  $\Rightarrow$  Clausius”:

- Clausius is wrong  $\Rightarrow$   $M_1$  possible.




# Equivalence



“Kelvin  $\Rightarrow$  Clausius”:

- Clausius is wrong  $\Rightarrow$   $M_1$  possible.
- For  $M_2$ ,  $\Rightarrow$   $W = Q_1 - Q$ .


# Equivalence



“Kelvin  $\Rightarrow$  Clausius”:

- Clausius is wrong  $\Rightarrow$   $M_1$  possible.
- For  $M_2$ ,  $\Rightarrow$   $W = Q_1 - Q$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1 - Q$  from  $T_1$ ;


# Equivalence



“Kelvin  $\Rightarrow$  Clausius”:

- Clausius is wrong  $\Rightarrow$   $M_1$  possible.
- For  $M_2$ ,  $\Rightarrow$   $W = Q_1 - Q$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1 - Q$  from  $T_1$ ;
- while to do  $W = Q_1 - Q$  to the outside;

# Equivalence



“Kelvin  $\Rightarrow$  Clausius”:

- Clausius is wrong  $\Rightarrow$   $M_1$  possible.
- For  $M_2$ ,  $\Rightarrow$   $W = Q_1 - Q$ .
- For  $M_1 + M_2$ , it is possible to absorb  $Q_1 - Q$  from  $T_1$ ;
- while to do  $W = Q_1 - Q$  to the outside;
- This contradicts with Kelvin statement.

## References

- F. Reif, *Fundamentals of Statistical And Thermal Physics*, 1st version.
- P. Jacobs, *Thermodynamics*, Imperial College Press, 2013.

# Acknowledgement

**Thank you for your attention!**